

Phosphatidylserine-dependent ingestion of apoptotic cells promotes TGF- β 1 secretion and the resolution of inflammation

Huynh, Fadok, Henson

The Journal of Clinical Investigation


2002

DOI:10.1172/JCI200211638






background

apoptosis
ancient Greek
'falling off'



- loss of PL asymmetry







© 2014 Pearson Education, Inc.

- embryogenesis & adult: maintain tissue f(x) & structure
- engulfed by (non-)professional phagocytes
- |-| release of potentially toxic / immunogenic content
- clinical implication



mφ that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF- β , PGE2, & PAF
 Fadok, et al., 1998, J Clin Invest



purpose of this study

TGF- β 1 & R expressed by almost all cells

-| growth & dx/dt

regulates immune & infl responses

modulates wound healing, ECM composition, adhesion & migration


apoptotic cell recognition systems: PSR, vitronectin R, CD14, collectin R

- determine the role of TGF- β 1 in clearance of apop cells
- elucidate a potential role of PS for TGF- β 1 production following ingestion of apop cells



m&m's

- human T lymphocyte Jurkat cell line
- human monomyelocyte PLB-985





results



TGF- β 1 production of inflamed sites-derived m φ



cell-associated TGF- β 1



DEPARTMENT OF OPHTHALMOLOGY
AND OPTOMETRY
MEDICAL UNIVERSITY OF VIENNA

J. Clin. Invest. 109:41-50 (2002).
DOI:10.1172/JCI2002638


ApoJ-induced TGF- β 1 secretion correlates w/ φcytic uptake



**ApoJ in non-inflamed
tissues failed to induce
TGF secretion**




TGF- β 1 secretion results from increased release & *de novo* synthesized TGF- β 1




apoptotic cell recognition is specific for TGF- β 1 induction


ms- α h CD45 (LCA)
IgG opsonization
Fc γ



Requirement of surface PS in apoptotic cell recognition for TGF- β 1 induction




Requirement of surface PS in apoptotic cell recognition for TGF- β 1 induction, c'd




ApoJ-induced reduction of pro-inflammatory mediators

Δ 1d post cell
instillation



mp inflammatory protein2
CXCL2
chemokine 4 leukocytes



cytokine-induced
neutrophil chemoattractant



monocyte chemoattractant
protein 1
CCL2
monocytes, memTcells, DC



ApoJ resulttet in reduction of infiltrated BALF mφ on d5




ApoJ result in reduction of infiltrated BALF mφ c'd


LPS

significant reduction in parenchymal infiltrates and intra-alveolar cells


a




b




Day 1 ApoJ




Day 3 ApoJ



Resolution of inflammation is specific for apoptotic cell clearance and is mediated by TGF



Resolution of inflammation is specific for apoptotic cell clearance and is mediated by TGF, c'd



concluding remarks

- instillation of apoptotic cells promotes TGF- β 1 production in inflamed tissues
- mφ-derived TGF- β 1 is preformed & *de novo* synthesized
- apoptotic cell recognition, but not Fcγ phagocytic system induces TGF secretion
- PS on apoptotic cell surface is crucial for TGF secretion
- reduction of pro-inflammatory mediators & infl cell infiltrates by instillation of apop cells
- **resolution of inflammation is specific for apoptotic cell clearance & mediated by TGF- β 1 & dependent on surface PS**



thank you

